



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2013

Superficial mycoses

Rostaher, A

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-81460>

Conference or Workshop Item

Originally published at:

Rostaher, A (2013). Superficial mycoses. In: Annual Meeting of the Slovene Dermatology Study Group, Bled, Slovenia, 29 June 2013.

Superficial mycoses

Dermatophytosis

Dermatophytosis is an infection of keratinized tissues (claws, hair and skin) usually caused by *Microsporum canis*, in some cases also by *M. gypsum* and *Trichophyton mentagrophytes*. The incidence and prevalence depends largely on the climatic condition and natural reservoir. In cats, *M. canis* is present all year round, in contrast *M. gypsum* and *T. mentagrophytes* are seldom reported in cats and are observed more frequently in the summer and fall months. The transmission is possible thru direct contact or indirectly (combs, brushes, transport cages, etc). The environment (hair shafts with fungal spores remain infectious for more then 18 months) and asymptomatic carries can act as reservoir for infection. Infections are more common in young, old and immunosuppressed animals. Additionally cats in catteries, shelters and pet shops can have a higher incidence of the disease.

Classical clinical lesions are annular alopecia, scaling, broken hair usually on the head, pinnae and paws. Other reported uncommon clinical presentations are miliary dermatitis, chin acne, kerion, onychomycosis, dermatophytic pseudomycetoma, otitis externa, eosinophilic plaques and recently described 'urticaria pigmentosa-like dermatitis. The diagnosis can be made by different techniques. Wood's lamp examination is positive in less then 50% of infected cats with *M. canis*. Direct examination of the hairs or even better skin scrapings offers a very good diagnostic tool. A fungal culture is required for confirmation of the diagnosis and for the determination of the fungal species.

The principles of a successful treatment are:

- Treat the affected cat (systemically and topically)
- Examination (fungal culture) and treatment of contact animals
- Decreasing environmental contamination (enilconazole, 1:10 diluted bleach)
- Stop treatment only after a negative fungal culture

Systemic medications for dermatophytosis include itraconazole (10 mg/kg daily or as pulse therapy). Other drugs reported to be efficacious are fluconazole, terbinafin and griseofulvin. The last one used less frequently due to the side effects such as panleukopenia, anorexia, vomiting... Especially, griseofulvin should be discouraged in FIV positive cats. Topical preparations are enilconazole, miconazole and lime sulfur and should be used twice weekly.

Clipping the cats can hasten the cure process, as it removes immediately a huge amount of spores, it helps to improve the efficacy of the topical treatment and also it leads to a decreased environmental contamination. There is still missing evidence of the efficacy of vaccines and lufenuron in the treatment of canine and feline dermatophytosis.

Malassezia dermatitis

Malassezia spp. are commensal yeasts that colonize the superficial epidermis of the skin and the ears. In cats the lipid non-dependend *M. pachydermaitis* and the lipid-non-dependent *M. sympodialis*, *M. globosa*, *M. nana*, *M. slooffiae*, *M. restricta* and *M. furfur* are usually found. *Malassezia* dermatitis is usually present due to an underlying disease, such as allergic dermatitis (of any origin), paraneoplastic syndrome (exfoliative dermatitis associated with thymoma, paraneoplastic alopecia) and FIV/FELV infection. Primary *Malassezia* dermatitis is rarely found in the cat. Hairless breeds or breeds with naturally reduced amounts of hairs such as the Sphynx

and the Rex cat are reported to carry a higher number of these fungi when compared to European short-haired breeds.

Predilection sites include the ear canals, interdigital spaces, ventral tail base, groin and areas with skin folds. The following clinical presentations associated with possible pruritus are usually seen in cats:

- Otitis externa with black and waxy otitis externa
- Chin acne
- Facial dermatitis
- Paronychia
- Generalized erythematous scaly to waxy dermatitis

Differential diagnoses include allergic dermatitis, superficial staphylococcal folliculitis, demodicosis, dermatophytosis, feline acne and epitheliotropic lymphoma.

The diagnosis is made by cytological examination using a microscopic slide, swab or a cellophane tape (for less accessible areas such as interdigital or dry lesions) and Diff-Quick stain. The fungal culture to diagnose *Malassezia* dermatitis is of no diagnostic value, as cats can be healthy carriers of these fungi.

Treatment options depend on the lesion type and area of distribution. Small focal areas can be treated with antifungal cream, wipes or ear drops (if the owner can prevent the cat to lick of the preparation min 5-10 minutes from the skin). Topical treatment in generalized disease included application of shampoos (containing chlorhexidin and miconazole) or enilconazole rinses. In the dog it was shown that the fastest clinical resolution is achieved with a combination of the topical and systemic treatment. This is often difficult to apply in the feline dermatology, as cats tend to hate topical products. Therefore, the best option is the utilization of a systemic antifungal such as itraconazole.

Ahman S, Perrins N, Bond R. Treatment of *Malassezia pachydermatis*-associated seborrheic dermatitis in Devon Rex cats with itraconazole--a pilot study. *Vet Dermatol.* 2007; 18(3): 171-4

Ahman SE, Bergström KE. Cutaneous carriage of *Malassezia* species in healthy and seborrheic Sphynx cats and a comparison to carriage in Devon Rex cats. *J Feline Med Surg.* 2009; 11(12): 970-6.

Bensignor E. Treatment of *Malassezia* overgrowth with itraconazole in 15 cats. *Vet Rec.* 2010 Dec 25;167(26):1011-2.

Bettenay S. How to effectively diagnose and manage dermatophytosis Proceedings of the 7th World Congress of Veterinary Dermatology, July 24-28 2012, Vancouver Canada, 9-17.

Bond R. Superficial veterinary mycoses. *Clin Dermatol.* 2010 Mar 4;28(2):226-36.

Carlotti DN, Guinot P, Meissonnier E, Germain PA. Eradication of feline dermatophytosis in a shelter: a field study. *Vet Dermatol.* 2010; 21(3): 259-66.

Colombo S, Nardoni S, Cornegliani L, Mancianti F Prevalence of *Malassezia* spp. yeasts in feline nail folds: a cytological and mycological study. *Vet Dermatol.* 2007 Aug;18(4):278-83.

Colombo S, Scarampella F, Ordeix L, Roccabianca P. Dermatophytosis and papular

eosinophilic/mastocytic dermatitis (urticaria pigmentosa-like dermatitis) in three Devon Rex cats. *Journal of Feline Medicine and Surgery* (2012) 14, 498–502.

Friberg C. Feline Facial Dermatoses. *Vet Clin North Am Small Anim Pract.* 2006 Jan; 36(1): 115-40

Forster-Van Hijfte MA, Curtis CF, White RN. Resolution of exfoliative dermatitis and *Malassezia pachydermatis* overgrowth in a cat after surgical thymoma resection. *J Small Anim Pract.* 1997; 38(10): 451-4.

Godfrey DR. A case of feline paraneoplastic alopecia with secondary *Malassezia*-associated dermatitis. *J Small Anim Pract.* 1998; 39(8): 394-6.

Jazic E, Coyner KS, Loeffler DG, Lewis TP. An evaluation of the clinical, cytological, infectious and histopathological features of feline acne. *Vet Dermatol.* 2006 Apr; 17(2):134-40.

Laura Ordeix, Franca Galeotti, Fabia Scarpella, Carla Dedola, Mar Bardagi, Erica Romano and Alessandra Fondati. *Malassezia* spp. overgrowth in allergic cats. *Vet Dermatol.* 2007; 18(5): 316-23.

Miller WH, Griffin CE, Campbell KL. Fungal and algal Skin Diseases. *Mueller and Kirk's Small Animal Dermatology*, 7th Edition, 2013, Elsevier, 223-283.

Moriello KA. Diagnostic techniques of dermatophytosis. *Clinical Techniques in Small Animal Practice* 2001; 16: 219–24

Moriello KA. Treatment of dermatophytosis in dogs and cats: review of published studies. *Vet Dermatol.* 2004; 15(2): 99-107.

Muller A, Guaguère E, Degorce-Rubiales F, Bourdoiseau G. Dermatophytosis due to *Microsporum persicolor*: a retrospective study of 16 cases. *Can Vet J.* 2011; 52(4): 385-8.

Nardoni S, Mancianti F, Rum A, Corazza M. Isolation of *Malassezia* species from healthy cats and cats with otitis. *J Feline Med Surg.* 2005 Jun;7(3):141-5.